



UDC 630.3

DOI: 10.48077/scihor.24(1).2021.77-84

ANALYSIS OF SPECIES DIVERSITY OF FORESTS ON THE TERRITORY OF KHARKIVSKA OBLAST

Oleksandr Bondar^{1*}, Liudmyla Tkach², Nelia Tsytsiura¹,
Oksana Halahan¹, Olena Tryhuba¹

¹Taras Shevchenko Regional Humanitarian-Pedagogical Academy of Kremenchuk
47003, 1 Litseina Str., Kremenchuk, Ukraine

²O.M. Beketov National University of Urban Economy in Kharkiv
61000, 17 Marshal Bazhanov Str., Kharkiv, Ukraine

Article's History:

Received: 01.12.2020

Revised: 14.01.2021

Accepted: 27.02.2021

Suggested Citation:

Bondar, O., Tkach, L., Tsytsiura, N., Halahan, O., & Tryhuba, O. (2021). Analysis of species diversity of forests on the territory of Kharkivska Oblast. *Scientific Horizons*, 24(1), 77-84.

Abstract. The paper presents the findings of a study of species diversity of forests in the Kharkivska Oblast. The study was conducted on the territory of ten forestry enterprises and the State Enterprise Kharkiv Forest Research Station and the Homilsha Woods National Nature Park on an area of 282.3 thousand ha (area is subordinated to the State Forest Resources Agency of Ukraine). The purpose of the study was to analyse the species diversity and typological structure of forests in the Kharkivska Oblast. The study of species diversity was carried out based on analysis of the subcompartment database of the VO Ukrderzhlisproekt. The plantations were classified by the trophotope on the territory of the research facility in accordance with the main methodological provisions of the Ukrainian forestry and forest typology. A diagrammatic map of forests and dominant species (common oak and Scots pine) was compiled using the MapInfo Professional 12.5 software package and a vector map of 12 forestries in the Kharkivska Oblast. It has been established that 56 species of trees grow in the Kharkivska Oblast. Among these tree species, the predominant ones are common oak (53.4%) and Scots pine (32.7%). The distribution of plantations by trophotope is characterised by certain features. For example, dubrava conditions prevail in the stands (65.6% of the total area covered by forest vegetation), subor forests are 3.5 times less represented, and sudubravas and pine forests are the least common. Among the forest types, the most common are fresh maple-linden dubravas (43.7%) and fresh oak-pine subors (18.9%). The practical significance of the study is that the analysis of species diversity and typological structure of forests must be taken into account during the planning, organisation, and conduct of forestry activities in the Kharkivska Oblast

Keywords: common oak, Scots pine, tree species, forest site types, trophotope, state enterprise, national nature park



Copyright © The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (<https://creativecommons.org/licenses/by/4.0/>)

*Corresponding author.

INTRODUCTION

The species diversity of forests is declining under the influence of human economic activity and climate change. This impact is global and irreversible [1]. There is a significant probability of changes in the most common species (Scots pine and common oak) of Ukraine, which in turn will lead to the transformation or even destruction of forest natural ecosystems that exist today [2]. If some decisive action is not taken now, further adaptation to climate change will require significant efforts and financial costs [3].

Forests for their intended purpose and location perform mainly water-conserving [4], protective [5], health-improving [6], aesthetic [7], educational and other functions [8]. Forests have sanitary and recreational value, cause a positive effect on climate, soil conditions, and water resources [9], reduce the effects of water [10] and wind erosion [11; 12], and is also a source of forest resources [13]. The formation and distribution of forest depends on the geomorphological structure, soil conditions, terrain and climatic features. All these factors must be taken into account during the planning, organisation, and management of forestry activities [14].

A significant number of papers, both by Ukrainian [15-17] and foreign [18-20] scientists, are devoted to the study of the species and typological structure of forests. In Ukraine, studies of the typological diversity of forests were mainly conducted within natural areas: Polissya, Forest-Steppe and Steppe. Regarding the study of species diversity of forests in the Kharkivska Oblast, recently such studies have been almost non-existent [21; 22]. The scientific novelty of the findings lies in the fact that the species and typological structure (by forest site types, trophotopes, and forest types) of forest plantations in Kharkivska Oblast was analysed for the first time, including diagrammatic maps of predominant species (Scots pine and common oak). The *purpose* of the study was to identify the features of the species diversity of forests in the Kharkivska Oblast.

To achieve the stated goal, the following research objectives were set:

- to analyse the distribution of plantations by tree species and forest types;
- to compile a map of forests and predominant species (Scots pine and common oak);
- to determine the number of tree species, forest site types, and trophotopes on the territory of forestry enterprises in Kharkivska Oblast.

MATERIALS AND METHODS

Species diversity of forests of state forestry enterprises located in the Kharkivska Oblast, which are subordinated to the State Forest Resources Agency of Ukraine, was determined based on the forest inventory materials as of 01.01.2016. The subcompartment databases of Ukrderzhlisproekt were converted using the NewUn-PackOHOTA software package, which was developed at the URIFFM named after G.M. Vysotsky [23].

Study was conducted in several stages. At the first stage, materials of forest inventory of ten state-owned enterprises (SE Balakliyske forestry, SE Blyzniukivske forestry, SE Vovchanske forestry, SE Hutyanske forestry, SE Zhovtneve forestry, SE Zmiivske forestry, SE Izyumske forestry, SE Krasnogradske forestry, SE Kupyanske forestry, SE Chuguevo-Babchanske forestry), Kharkiv Forest Research Station, and Homilsha Woods National Nature Park were put together and analysed (Fig. 1). These materials (databases) were grouped by certain indicators. The filtering of information was carried out in the following sequence: tree species, forest site types, trophotopes, and forest types. The analysis of these data was performed using appropriate computer software (*Microsoft Excel 2010*), as well as geoinformation technologies (*MapInfo Professional 12.5. Beta*). The silvicultural and typological analysis of forests was carried out in accordance with the main methodological provisions of the forestry and ecological (Ukrainian) consensus group of forest typology [24].




Figure 1. Diagrammatic map of forest distribution in the Kharkivska Oblast

At the second stage, diagrammatic map of forests, distribution of predominant tree species (common oak and Scots pine) in the Kharkivska Oblast has been compiled. It was done using *MapInfo Professional 12.5. Beta* and vector map of Ukraine. Thus, the geographical coordinates of the extreme points of the survey plot: in the North – 37° 29'33", 50° 27'26", in the South – 36° 17'50", 48° 31'56", in the West – 34° 51'27", 49° 59'11", and in the East – 38° 05'36", 49° 50'43". The total area of the investigated forests, which are subordinated to the State Forest Resources Agency of Ukraine, is 282.3 thousand hectares. The distribution of forests in the Kharkivska Oblast is uneven. They are mostly located in the floodplain of the Siverskyi Donets river and its tributaries.

RESULTS AND DISCUSSION

Species composition of forests in the Kharkivska Oblast is quite diverse. Thus, the largest concentration of species is in SE Izyumske forestry (37 species of trees) and SE Zhovtneve forestry (36 species of trees), and the least – in SE Chuguevo-Babchanske forestry (21 species of trees) and the Homilsha Woods NNP (16 species of trees), respectively (Table 1). A total of 18 forest site types have been identified on the territory of research facility. Thus, most of them are concentrated in the Izyumske forestry (18 site types), and the least – in the Homilsha Woods NNP (8 site types).

The distribution of forests in the Kharkivska Oblast

by trophotopes is characterised by certain peculiarities. Thus, the plantations are dominated by dubrava site type (65.6% of the total area covered by forest vegetation), 3.5 times less represented are subors (18.3%), and the least represented are sudubravas (9.9%) and pine forests 6.2%).

In total, 18 types of forest sites have been identified in the forests of the Kharkivska Oblast, in particular, in pine forests there are 3 site types, and in subors, sudubravas and dubravas – 5 site types in each. Pine forests with poor growing conditions are present on the territory of 10 enterprises, and their share varies from 0.3% (SE Kharkiv Forest Research Station) to 27.7% (SE Izyumske forestry). On the territory of two enterprises (SE Blyzniukivske forestry and Homilsha Woods NNP) pine woods are not detected at all. Subor types are common on the territory of all enterprises, and their share varies from 0.3% (Homilsha Woods NNP) to 32.3% (SE Izyumske forestry). Sudubrava types are present on the territory of all enterprises, and their share ranges from 2.1% (SE Kharkiv Forest Research Station) to 25.4% (SE Hutyanske forestry). In dubrava (rich) conditions, plantations grow on the territory of all enterprises, and their share varies from 25.1% (SE Izyumske forestry) to 94.4% (Homilsha Woods NNP). After analysing the distribution of plantating by trophotopes in the Kharkivska Oblast (Fig. 2), it was determined that the natural conditions are favourable for growing high-yielding plantations of common oak and Scots pine.

Table 1. Distribution of forest areas in the Kharkivska Oblast

State-owned enterprise	Number		Area	
	tree species	forest site types	ha	%
SE Balakliyske forestry	29	17	22,650.0	8.0
SE Blyzniukivske forestry	25	11	5,350.8	1.9
SE Vovchanske forestry	27	16	25,830.0	9.1
SE Hutyanske forestry	30	16	27,686.1	9.8
SE Kharkiv Forest Research Station	33	14	19,753.7	7.0
SE Zhovtneve forestry	36	16	45,321.8	16.1
SE Zmiivske forestry	28	17	23,752.1	8.4
SE Izyumske forestry	37	18	43,101.8	15.3
SE Krasnogradske forestry	28	15	12,083.3	4.3
SE Kupyanske forestry	33	18	33,633.1	11.9
Homilsha Woods NNP	16	8	3167.7	1.1
SE Chuguevo-Babchanske forestry	21	15	19,999.0	7.1
Total	56	18	282,329.4	100.0




Figure 2. Distribution of plantation areas of enterprises by trophotypes, %

In each type of forest sites, foresters have allocated a different number of forest types. The pine forests are represented by four, the subors by seven, the sudubravas by eleven, and the dubravas by thirteen forest types. A total of 35 forest types were identified: one was very dry, six were dry, ten were fresh, nine were moist, five

were wet, and four were damp (Table 2). The typological variety of forests in the Kharkivska Oblast is quite diverse. Thus, the most common type of forest is fresh maple-linden dubrava (43.9% of the total area covered with forest vegetation), a slightly smaller share is fresh oak-pine subor (18.9%) and dry maple-linden dubrava (11.9%).

Table 2. Forest types in the Kharkivska Oblast

Forest type		Area	
Name	Index	ha	%
Very dry pine forest	A ₀ -P	19.2	0.01
Dry pine forest	A ₁ -P	5,543.4	2.0
Fresh pine forest	A ₂ -P	17,131	6.1
Moist pine forest	A ₃ -P	11.5	0.0
Dry oak-pine subor	B ₁ -oP	5,243.2	1.9
Dry eroded oak subor	B ₁ -Oe	4.9	0.0
Fresh oak-pine subor	B₂-oP	53,285.8	18.9
Moist oak-pine subor	B ₃ -oP	1,348.9	0.5
Moist floodplain sedge subor	B ₃ -Sf	201.1	0.1
Wet oak-pine subor	B ₄ -oP	142.8	0.1
Damp birch-pine subor	B ₅ -bP	4.3	0.002
Dry eroded Tatarian maple sudubrava	C ₁ -TmOe	3,602.1	1.3
Fresh oak sudubrava	C ₂ -O	919.3	0.3
Fresh eroded linden sudubrava	C ₂ -lOe	2,072.6	0.7
Fresh linden-oak-pine sudubrava	C ₂ -loP	15,571.8	5.5
Fresh floodplain oak sudubrava	C ₂ -Of	1,525.8	0.5
Moist maple-linden sudubrava	C ₃ -mLO	287.8	0.1

Continuation of table 2

Forest type		Area	
Name	Index	ha	%
Moist linden-oak-pine sudubrava	C ₃ -loP	1,330.0	0.5
Moist floodplain oak sudubrava	C ₃ -Of	1,619.8	0.6
Wet alder sudubrava	C ₄ -A	2,062.7	0.7
Wet poplar-willow sudubrava	C ₄ -pW	236.5	0.1
Damp alder sudubrava	C ₅ -A	155.0	0.1
Dry elm-Tatarian maple dubrava	D ₁ -eTmO	711.0	0.3
Dry maple-linden dubrava	D₁-mLO	33,608.6	11.9
Fresh elm-Tatarian maple dubrava	D ₂ -eTmO	404.1	0.1
Fresh maple-linden dubrava	D₂-mLO	123239.2	43.7
Fresh ash-linden dubrava	D ₂ -alO	103.0	0.04
Fresh floodplain elm-Tatarian maple dubrava	D ₂ -etmOf	1,042.5	0.4
Moist maple-linden dubrava	D ₃ -mLO	707.9	0.3
Moist ash-linden dubrava	D ₃ -alO	1,144	0.4
Moist floodplain elm-Tatarian maple dubrava	D ₃ -etmOf	7,204.4	2.6
Wet alder dubrava	D ₄ -A	1,110.3	0.4
Wet floodplain poplar-willow dubrava	D ₄ -pWf	177.8	0.1
Damp alder dubrava	D ₅ -A	518.9	0.2
Damp floodplain willow dubrava	D ₅ -Wf	38.2	0.01
Total			100.0

The greatest typological diversity is characteristic of the forests of the State Enterprise Izyumske forestry (34 forest types), and the least – for the forests of the Homilsha Woods NNP (12 forest types). The small number

of forest types is explained by the homogeneity of geomorphological, soil, and climatic conditions of plantation within the relevant forestry enterprises (Fig. 3).




Figure 3. Distribution of the total number of forest types by enterprises

The nature of the distribution of wood species on the territory of the research facility is quite uneven (Fig. 4). Thus, in the Kharkivska Oblast there are 56 species of trees, of which the most common (Table 3) are common

oak (*Quercus robur* L.) – 53.4% (150798.4 ha of the total area covered by forest vegetation) and Scots pine (*Pinus sylvestris* L.) – 32.7% (92380.0 ha).

Table 3. Distribution of areas of predominant tree species in the Kharkivska Oblast, ha*

Tree species	Total	
	ha	%
European birch	2,727.1	1.0
Common alder	4,404.9	1.6
European elm	513.2	0.2
Common oak	150,798.4	53.4
Norway maple	3,387.0	1.2
Tatarian maple	1,248.9	0.4
Big-leaf linden	1,662.9	0.6
Black locust tree (false acacia)	7,142.3	2.5
Scots pine	92,380.0	32.7
European aspen	3,997.6	1.4
Silver poplar	1,622.5	0.6
Canadian poplar	752.9	0.3
Black poplar	684.2	0.2
European ash	7,062.3	2.5
Green ash	1,640.6	0.6
Other	2,304.6	0.8
Total	282,329.4	100.0

Note: * The table shows the data of the predominant tree species, the area of which is more than 150 ha, from the total area covered with forest vegetation on the territory of research facility




Figure 4. Diagrammatic map of the distribution of dominant tree species in the Kharkivska Oblast

With the increase of areas covered by forest vegetation, the number of forest types and tree species increases. This complicates the conditions for forest management activities in Kharkivska Oblast, and, in turn, increases biological diversity.

CONCLUSIONS

The species diversity of forests on the territory of the research facility is represented by 56 tree species. Among them, the largest share is occupied by common oak (53.4% of the total area covered with forest vegetation) and Scots pine (32.7%). Forest management activities on the territory of the research facility should be aimed at increasing the species diversity of forests by carrying

out certain economic activities.

The trophotopes are dominated by dubrava (oak) conditions, the share of which is 65.6% of the total area covered with forest vegetation. This contributes to the cultivation of highly productive oak forests. The typological structure of forests is represented by 35 forest types. Among the forest types, the most predominant are fresh maple-linden dubravas (43.9% of the total area covered with forest vegetation), fresh oak-pine subors (18.9%) and dry maple-linden dubravas (11.9%). According to the compiled diagrammatic maps of forests it is possible to determine the actual amount of woodland and peculiarities of plantation distribution. These materials can be used when creating new forests.

REFERENCES

- [1] Case, M.J., Johnson, B.G., Bartowitz, K.J., & Hudiburg, T.W. (2021). Forests of the future: Climate change impacts and implications for carbon storage in the Pacific Northwest, USA. *Forest Ecology and Management*, 482, article number 118886. doi: 10.1016/j.foreco.2020.118886.
- [2] Ivanyuta, S.P., Kolomiets, O.O., Malinovskaya, O.A., & Yakushenko, L.M. (2020). *Climate change: Consequences and adaptation measures*. Kyiv: NISD.
- [3] Viccaro, M., Cozzi, M., Fanelli, L., & Romano, S. (2019). Spatial modelling approach to evaluate the economic impacts of climate change on forests at a local scale. *Ecological Indicators*, 106, article number 105523.
- [4] Butzen, V., Seeger, M., Marruedo, A., Jonge, L., Wengel, R., Ries J., & Casper, M. (2015). Water repellency under coniferous and deciduous forest – Experimental assessment and impact on overland flow. *Catena*, 133, 255-265. doi: 10.1016/j.catena.2015.05.022.
- [5] Miura, S., Amacher, M., Hofer, T., San-Miguel-Ayanz, J., & Thackway, R. (2015). Protective functions and ecosystem services of global forests in the past quarter-century. *Forest Ecology and Management*, 352, 35-46. doi: 10.1016/j.foreco.2015.03.039.
- [6] Rosa, C.D., Larson, L.R., Collado, S., & Profice, C.C. (2021). Forest therapy can prevent and treat depression: Evidence from meta-analyses. *Urban Forestry & Urban Greening*, 57, 126943. doi: 10.1016/j.ufug.2020.126943.
- [7] Urbis, A., Povilanskas, R., & Newton, A. (2019). Valuation of aesthetic ecosystem services of protected coastal dunes and forests. *Ocean & Coastal Management*, 179, article number 104832. doi: 10.1016/j.ocecoaman.2019.104832.
- [8] Glatthorn, J., Annighöfer, P., Balkenhol, N., Leuschner, C., Polle, A., Scheu, S., Schuldt, A., Schuldt, B., & Ammer, C. (2021). An interdisciplinary framework to describe and evaluate the functioning of forest ecosystems. *Basic and Applied Ecology*, 52, 1-14. doi: 10.1016/j.baae.2021.02.006.
- [9] Luo, P., Zhou, M., Deng, H., Lyu, J., Cao, W., Takara, K., Nover, D., & Schladow, S. (2018). Impact of forest maintenance on water shortages: Hydrologic modeling and effects of climate change. *Science of The Total Environment*, 615, 1355-1363. doi: 10.1016/j.scitotenv.2017.09.044.
- [10] Teng, M., Huang, C., Wang P., Zeng, L., Zhou, Z., Xiao, W., Huang, Z., & Liu, C. (2019). Impacts of forest restoration on soil erosion in the Three Gorges Reservoir area, China. *Science of The Total Environment*, 697, article number 134164. doi: 10.1016/j.scitotenv.2019.134164.
- [11] Guo, B., Zang, W., Yang, X., Huang, X., Zhang, R., Wu, H., Yang, L., Wang, Z., Sun, G., & Zhang, Y. (2020). Improved evaluation method of the soil wind erosion intensity based on the cloud-AHP model under the stress of global climate change. *Science of The Total Environment*, 746, article number 141271. doi: 10.1016/j.scitotenv.2020.141271.
- [12] Zöll, U., Lucas-Moffat, A.M., Wintjen, P., Schrader, F., Beudert, B., & Brümmer, C. (2019). Is the biosphere-atmosphere exchange of total reactive nitrogen above forest driven by the same factors as carbon dioxide? An analysis using artificial neural networks. *Atmospheric Environment*, 206, 108-118. doi: 10.1016/j.atmosenv.2019.02.042.
- [13] Freundorfer, A., Rehberg, I., Law, B.E., & Thomas, C. (2019). Forest wind regimes and their implications on cross-canopy coupling. *Agricultural and Forest Meteorology*, 279, article number 107696. doi: 10.1016/j.agrformet.2019.107696.
- [14] Tkach, V.P. (1999). *Floodplain forest of Ukraine*. Kharkiv: Pravo.
- [15] Siruk, Yu.V., Pechenyuk, E.P., & Chernyuk, T.N. (2015). Typological structure and characteristics of forest fund of the Central Polyssya of Ukraine. *Scientific Bulletin of UNFU*, 25(10), 97-103. doi: 10.15421/40251014.
- [16] Kratiuk, O.L. (2019). Typological structure of forest plantations and biotopic characteristics of enclosures in Central Polissya. *Scientific Bulletin of UNFU*, 29(2), 62-64. doi: 10.15421/40290212.

- [17] Muradyan, L.V., & Sytnyk, S.A. (2020). Typological structure and valuation features of black locust stands in protective forests within the Ravine Steppe of Ukraine. *Scientific Bulletin of UNFU*, 30(5), 36-41. doi: 10.36930/40300506.
- [18] Qian, J., Zhuang, H., Yang, W., Chen, Y., Chen, S., Qu, Y., Zhang, Y., Yang, Y., & Wang, Y. (2020). Selecting flagship species to solve a biodiversity conservation conundrum. *Plant Diversity*, 42(6), 488-491. doi: 10.1016/j.pld.2021.01.004.
- [19] Tarabon, S., Dutoit, T., & Isselin-Nondedeu, F. (2021). Pooling biodiversity offsets to improve habitat connectivity and species conservation. *Journal of Environmental Management*, 277, article number 111425. doi: 10.1016/j.jenvman.2020.111425.
- [20] Saraev, V., Valatin, G., Peace, A., & Quine, C. (2019). How does a biodiversity value impact upon optimal rotation length? An investigation using species richness and forest stand age. *Forest Policy and Economics*, 107, article number 101927. doi: 10.1016/j.forpol.2019.05.013.
- [21] Musienko, S., Lyalin, O., Tkach, L., Bondarenko, V., Kolenkina, M., & Kolchanova, O. (2020). State and structure of urban forests in Kharkivska Oblast. *Folia Forestalia Polonica, Series A – Forestry*, 62(1), 46-52. doi: 10.2478/ffp-2020-0006.
- [22] Bondar, O., Rumiantsev, M., Tkach, L., & Obolonyk, I. (2020). Prevailing forest types in the river catchments within the Left-Bank Forest-Steppe zone, Ukraine. *Folia Forestalia Polonica, Series A – Forestry*, 62(2), 100-113. doi: 10.2478/ffp-2020-0011.
- [23] Ostapenko, B.F., & Tkach, V.P. (2002). *Forest typology*. Kharkiv: KhNAU named after V.V. Dokuchaieva.
- [24] Vedmid, M.M., Mieshkova, V.L., & Zhezhkun, A.M. (2006). Algorithm for revealing the land of youngsters in dibrovah for materials lisovorderkuvannya. *Forestry and Forest Melioration*, 110, 54-59.

АНАЛІЗ ВИДОВОГО РІЗНОМАНІТТЯ ЛІСІВ НА ТЕРИТОРІЇ ХАРКІВСЬКОЇ ОБЛАСТІ

Олександр Богданович Бондар¹, Людмила Іванівна Ткач², Неля Іванівна Цицюра¹,
Оксана Костянтинівна Галаган¹, Олена Василівна Тригуба¹

¹Кременецька обласна гуманітарно-педагогічна академія ім. Тараса Шевченка
47003, вул. Лицейна, 1, м. Кременець, Україна

²Харківський національний університет міського господарства ім. О.М. Бекетова
61000, вул. Маршала Бажанова, 17, м. Харків, Україна

Анотація. Наведено результати досліджень видового різноманіття лісів на території Харківської області. Дослідження проведено на території десяти лісогосподарських підприємств, Харківської лісової науково-дослідної станції та Національного природного парку «Гомільшанські ліси» на площі 282,3 тис. га, які знаходяться у підпорядкуванні Держлісагенства України. Метою дослідження було проаналізувати видове різноманіття та типологічну структуру лісів на території Харківської області. Дослідження видового різноманіття здійснювали на основі аналізу повидільної бази даних ВО «Укрдержліспроєкт». Класифікацію насаджень за трофотопами на території дослідного об'єкта здійснювали згідно із основними методичними положеннями лісівничо-екологічної (української) школи лісової типології. Побудовано загальну схему-карту лісів і переважаючих порід (дуба звичайного та сосни звичайної) на основі програми *MapInfo Professional 12.5* і векторної карти 12 дослідних об'єктів на території Харківської області. Визначено, що на території Харківської області ростуть 56 видів деревних порід. Серед деревних порід найбільш поширеними є дуб звичайний (53,4 %) та сосна звичайна (32,7 %). Розподіл насаджень за трофотопами характеризується певними особливостями. Так, у насадженнях домінують грудові умови (65,6 % від загальної площі вкритої лісовою рослинністю земель), у 3,5 рази менше представлені субори (18,3 %), а найменш представлені є сугруди (9,9 %) і бори (6,2 %). Серед типів лісу найбільш поширеним є свіжа кленово-липова діброва (43,7 %) і свіжий дубово-сосновий субір (18,9 %). Практична цінність статті полягає в тому, що результати аналізу видового різноманіття і типологічної структури лісів необхідно враховувати під час планування, організації та проведенні лісогосподарських заходів на території Харківської області

Ключові слова: дуб звичайний, сосна звичайна, деревна порода, типи лісорослинних умов, трофотоп, державне підприємство, національний природний парк